Fax: 949-		Die Scierosis Enrollment Form340-8008Phone: 949-305-0788Request			
Whole Health Pharmacy Rep:					
PATIENT INFORMA	ATION	PRESCRIBER INFORMATION			
Please complete the following or send patient demographi c sheet Patient Name Address		Prescriber Name	Prescriber Name		
Address 2		NDT	DEA NPI		
City, State, Zip		Address	Address		
City, State, Zip Home PhoneMobile Phone DOBLast Four of SSGender		City, State, Zip	City, State, Zip PhoneFax		
Language Preference			PhoneFax Contact Person		
Clinical Diagnosis:	please fax or email relevant clinical	notes, labs, tests, and previous medical history to expedite pr	rior authorizatio	n	
Diagnosis - Please incl	ude diagnosis name with ICD-10 code	Additional information Therapy New Reauthorizatio	Additional information Therapy New Reauthorization Restart		
 G35 Multiple Sclerosis Other Diagnosis: ICD-10 Code 			🗆 Extavia 🗆 Tecfidera 🗆 Other		
Description Number of Relapses in Past Year		Treatment Response	Treatment Response Treatment Dates		
Date of Diagnosis		Allergies	Allergies		
Date of Diagnosis		No Lab Data	Lab Data		
Date of Screen:	Positive Negative (Required for Ocrevus)	Additional Comments	Concomitant Medications		
	PRES	SCRIPTION INFORMATION			
Medication	Dose/Strength	Directions	Quantity	Refills	
Aubagio	□ 7mg Tablet □ 14mg Tablet	Take 1 tablet by mouth once daily	 28 days 30 days 		
Betaseron	□ 0.3mg Vial & Diluent	 Inject 0.0625mg (0.25ml) SQ every other day and increase over a six-week period to 0.25mg (1ml) SQ every other day (Starter Dose) 	□ 28 days		
🗆 Copaxone	 20mg Syringe 40mg Syringe 	 İnject 0.25mg (1ml) SQ every other day Inject 20mg SQ once daily Inject 40mg SQ three times per week and at least 48 hours apart 	□ 28 days		
🗆 Glatopa	□ 20mg Syringe	Inject 20mg SQ once daily	□ 28 days		
🗆 Kesimpta	 40mg Syringe 20 mg/0.4 mL solution in a single- 	 Inject 40mg SQ three times per week and at least 48 hours apart Initial Dosing: 20 mg administered at Week 0, 1, and 2. 	□ 28 days		
	 dose prefilled Sensoready ® Pen 20 mg/0.4 mL solution in a single- dose PFS 	 Initial Dosing: 20 mg administered at week 0, 1, and 2. Subsequent Dosing: 20 mg administered monthly starting at Week 4 			
Mayzent	 0.25mg Tablet 2mg Tablet 	 Titration Dosing as directed Take 1mg by mouth once daily Take 2mg by mouth once daily 			
	□ 300mg/10 mL single dose vial	 300 mg intravenous infusion followed two weeks later by a second 300mg intravenous infusion (starter dose) 	□ 2 vials		
□ Solu-Medrol	□ 125mg Act-O-Vial System	 G00 mg intravenous infusion every 6 months 125 mg administered intravenously 30 minutes prior to each Ocrevus infusion 	□ 1 vial □ 2 vials		
diphenhydramine	□ 50mg/ml vial	 50mg administered intravenously 30-60 minutes prior to each Ocrevus infusion 	 1 vial 2 vials 		
Tecfidera	 120mg Capsule 240mg Capsule Starter Pack 	 Starting dose: 120 mg twice a day, orally, for 7 days Maintenance dose after 7 days: 240 mg twice a day, orally Take starter pack as directed 	□ 7 days □ 30 days		
Tysabri	□ 300mg/15 mL vial	300 mg intravenous infusion over 1 hour every 4 weeks	🗆 1 vial		
□ Solu-Medrol	□ 125mg Act-O-Vial System	125 mg administered intravenously 30 minutes prior to each Tysabri infusion	🗆 1 vial		
diphenhydramine	□ 50mg/ml vial	 50mg administered intravenously 30-60 minutes prior to each Tysabri infusion 	🗆 1 vial		
Zeposia	□ 7 day Starter Pack	Days 1-4 Take 0.23 mg once daily	1 Starter		
	 Starter Kit (1 month supply) 0.92 mg maintenance 	Days 5-7 Take 0.46 mg once daily Day 8 and thereafter Take 0.92mg once daily Take 1 capsule by mouth once daily	Pack □ 30 days		
Ship to: □ Patient □ Office □ OtherDate:Need by Date					
*Prescriber Authorization: I authorize this pharmacy and its representatives to act as my authorized agent to secure coverage and initiate the insurance prior authorization process for my patient(s) and to sign any necessary forms on my behalf as my authorized agent, including the receipt of any required prior authorizations forms and the receipt and submission of patient tab values and other patient data, in the event that this pharmacy determines that it is unable to fulfill this prescription, I further authorize the pharmacy to forward this information and any related materials related to coverage of the product to another pharmacy of the patient's choice or in the patient's insurer's provider network.					

Prescriber's Signature:

CONFIDENTIALITY STATEMENT: This communication is intended for use of the individual or entity to which it is addressed and may contain information that is privileged, confidential, and exempt from disclosure under applicable law If the reader of this communication is not the intended recipient or the employee or agent responsible for delivery of the communication, you are hereby notified that any dissemination distribution, or copying of the communication is strictly prohibited. If you have received this communication in error, please notify us immediately by telephone.

Date: